

(1991) American Journal of Medical Science 301:32-43). Congestive heart failure or death from cardiac arrhythmias is common (Koren, A. et al. (1987) Am. J. Dis. Child. 141:93-96). For disorders such as thalassemia major (by definition, transfusion-dependent thalassemia), iron-overload now is the limiting factor in survival. The advent
 5 of chronic transfusion therapy in the 1960's increased life span to the early to mid-twenties, however, the pernicious consequences of iron-overload were invariably fatal (Cooley, T. (1945) Am. J. Med. 209:561-572), (Piomelli, S. (1991) Hematol. Oncol. Clin. North. Am. 5:557-569).

Iron is one of the leading causes of pediatric poisoning deaths in the
 10 United States (Litovitz, T.L. et al. (1992) Am. J. Emerg. Med. 10:452-505). Numerous reports of serious or fatal poisonings have been cited in the medical literature (Litovitz, T.L. et al. (1992); Westlin, W.F. (1966) Clin. Pediatr. 5:531-535; Henriksson, P. et al. (1979) Scand. J. Haematol. 22:235-240) including five toddler deaths in Los Angeles county during a seven month period in 1992 (Weiss, B. et al. (1993) Morb. Mortal.
 15 Wkly. Rep. 42:111-113). It is clear that iron can cause serious morbidity and mortality, yet many clinicians and families remain unaware of the dangers of iron (Anderson, B.D. (April 18, 2000) Medscape Pharamcists). Although uncommon, iron solutions may be absorbed through damaged or burned skin. Following ingestion of large amounts of iron, peak serum levels generally occur within 2 to 6 hours. After ingestion, iron in the
 20 +2 state is oxidized to the +3 state and attached to the transport protein, ferritin. The iron is then released from the ferritin to transferrin in the plasma, transported to the blood forming storage sites, and incorporated into enzymes in the body. Iron is eliminated slowly from the body. Even in states of iron overload, children may lose up to 2 mg per day. Ingestion of less than 20 mg/kg elemental iron is likely to produce GI
 25 symptoms. For patients who ingest greater than 60 mg/kg elemental iron, potentially life threatening symptoms may occur.

Furthermore, the emergence of drug resistant parasites, e.g., malaria, has intensified the search for new therapeutic approaches (e.g. drug combinations). One new approach under investigation is the administration of iron chelating agents
 30 (Cabantchik, Z.I. et al. (1996) Acta Haematol. 95:70-77; Van Zyl, R.L. et al. (1992) J. Antimicrob. Chemother. 30:273-278).

Patients with transfusion iron-overload, iron poisoning, and drug resistant parasitic diseases (e.g., malaria) are commonly treated with low molecular weight iron chelators. These compounds remove the excess, toxic iron from the patient's blood.
 35 The most commonly used drug worldwide is Desferrioxamine (Desferal®, Novartis). Therapy with Desferrioxamine is effective, but under-utilized because of drug delivery problems. Oral absorption of Desferrioxamine is very low. In some cases, Desferrioxamine infusion has proven not to be adequate (Westlin, W. (1996) Clin.

Ped.5:531-535; Tenenbein, M. et al. (1992) Lancet 339:699-701; Adamson, I.Y. et al. (1993) Toxicol. Appl. Pharmacol. 120:13-19). In addition, the low molecular weight of this hydrophilic molecule (657 Da) leads to renal clearance in about 15 to 20 minutes. Consequently, the drug is given by continuous infusion over 12 to 16 hours. This is
5 done either by subcutaneous infusion or by infusion into a permanent catheter. Such a long infusion duration is inconvenient and prone to infections and thrombosis. Desferrioxamine also has severe drawbacks in the treatment of parasitic diseases; (1) it is hydrophilic and poorly absorbed after oral administration; and (2) it is cleared rapidly after intravenous administration and iron chelators like Desferrioxamine do not readily
10 penetrate into advanced growth stages of parasitized cells (Loyevsky, M. et al. (1993) J. Clin. Invest. 91:218-224). As a consequence, continuous infusion of iron chelators like Desferrioxamine over a three day period is required to obtain enhanced parasite clearance in human malaria (Mabeza, G.F. et al. (1996) Acta Haematol. 95:78-86). Nonetheless, many patients use Desferrioxamine suboptimally or not at all.

15 No other chelator has proven clinical efficacy. Searches for clinically effective alternatives to Desferrioxamine for transfusional iron-overload have thus far been futile. Some chelating agents, such as diethyltriamine pentaacetic acid (DPTA) are effective, but too toxic for clinical use. Other chelators (e.g., EDTA) bind other cations in addition to iron, making them unacceptable as treatment of transfusional iron-
20 overload or iron poisoning.

One approach to the problem has been to immobilize Desferrioxamine to a large molecular matrix, thereby extending its biological half life. Immobilized Desferrioxamine depends on a shift in “pseudoequilibrium” conditions to produce a net outflux of iron from cells. The vast amount of storage iron exists inside cells, however,
25 effectively out of the reach of immobilized Desferrioxamine. The problem is that the storage iron inside the cells remains a dangerous source of free radicals until it is chelated and inactivated by the Desferrioxamine in the matrix.

The only chelator currently in extensive clinical trial is Deferipone (L1). Deferipone removes excess iron reasonably well although it falls short of
30 Desferrioxamine in this regard (Collins, A. et al. (1994) Blood 83:2329-33). The great appeal of Deferipone over Desferrioxamine is its oral absorption. For many patients, the convenience of an orally active chelator might more than compensate for lesser efficacy.

A number of clinical problems cloud Deferipone’s future. Severe agranulocytosis occurs in about 2% of patients (al-Refaie, F. et al. (1992) Blood 80:593-
35 9). Other significant side-effects of Deferipone include arthralgias and severe nausea (al-Refaie, F. et al. (1995) Br. J. Haematol. 91:224-229). Because of these and other problems, Deferipone’s clinical future is far from assured. Therefore, there exists a need for an improved iron chelator delivery system.

Summary of the Invention

The invention features an iron chelator delivery system comprising an iron chelator and a lipid carrier. The term "lipid carrier," as used herein, includes a carrier comprising lipid molecules, e.g., a liposome. The iron chelator and the lipid carrier of the iron chelator delivery system of the present invention can be combined in various ways. For example, the iron chelator can be coated with the lipids of the lipid carrier, e.g., iron chelators associated or attached to the surface of the lamellae of a lipid molecule or a liposome. Alternatively, the iron chelator can be encapsulated within the lipid carrier, e.g., iron chelators encapsulated within the central cavity of a liposome, intercalated within the lamellae of a liposome, or located between lamellae of the liposome.

Advantages of the iron chelator delivery system of the present application include: (1) a reduced toxicity based on the administration of a lower dose for a shorter period of time; (2) targeted delivery of the iron chelator without high renal clearance; (3) elimination of the pain associated with subcutaneous injections; (4) an increase in the half life of the iron chelator via targeted delivery of the drug to the heart and the liver, thus, reducing the amount of drug needed; and (5) entrapment of the drug in the liver up to 5 days, thus, allowing a longer period of time for iron chelation. Some iron chelation from the liver parenchyma will redistribute to other tissues and bind free iron.

Iron chelators within the scope of the present invention include, for example, Desferrioxamine, Deferipone, PIH (pyridoxal isonicotinoyl hydrazone and analogues), Rhodotorulic acid, HBED (N,N'-Bis(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid), HBPD (N,N'-Bis(2-hydroxybenzyl)propylene-1,3-diamine-N,N'-diacetic acid), 2,3-dihydroxybenzoic acid, DTPA [diethyltriamine pentaacetic acid] and iron chelators produced by bacterial siderophores. Lipid carriers within the scope of the present invention include, for example, liposomes, e.g., multilamellar and unilamellar liposomes, as well as, phospholipid and nonphospholipid liposomes.

In one embodiment, the concentration of the iron chelator within the iron chelator delivery system is approximately 1 μ M to 100mM. In another embodiment, the size of the liposome is approximately less than about 10 nanometers to about 10 microns. In a preferred embodiment, the iron chelator delivery system is targeted for delivery to the heart and liver. For example, the lipid carrier can comprise cationic or anionic charge groups or antibodies specific for cardiac proteins, e.g., myosin, troponin, or myosin light chain proteins, vasculature proteins, endothelial cells, or matrix proteins. In another preferred embodiment, the lipid carrier is galactosylated or mannosylated. In another preferred delivery system the iron chelator delivery system is tagged with a label

e.g. a radiolabel, which can be used for diagnostic imaging or identification of delivery to a particular site.

The present invention is also drawn to methods for preparing iron chelator delivery systems. Such methods comprise the steps of combining a lipid carrier or liposome with an iron chelator, and extracting the iron chelator-encapsulated liposomes to form an iron chelator delivery system.

Methods for treating iron-overload in a mammal, comprising administering to the mammal an iron chelator delivery system, e.g., an iron chelator associated with a lipid carrier, e.g., a liposome, are also provided. In a preferred embodiment, the iron chelator delivery system is administered so as to target specific organs, for example, the heart or the liver. Prior to administration, the iron chelator drug delivery system can be dissolved in a pharmaceutically acceptable excipient or carrier, e.g., saline, or water.

15 **Brief Description of the Drawings**

Figures 1 and 2 show the UV spectrum data for Desferrioxamine.

Figures 3 and 4 show the calibration curve for Desferrioxamine in methanol.

Figure 5 shows the blood chemistry values of normal animals before and after liposome infusion with and without Desferrioxamine.

Figures 6-18 show ^{111}In -Desferrioxamine biodistribution in rats.

Detailed Description of the Invention

The present invention provides an iron chelator delivery system comprising an iron chelator and a lipid carrier. Iron chelators are compounds that remove the excess, toxic iron from the patient's blood, tissues and organs and include, for example, Desferrioxamine, Deferipone, PIH (pyridoxal isonicotinoyl hydrazone), Rhodotorulic acid, HBED (N,N'-Bis(2-hydroxybenzyl)ethylenediamine-N,N-diacetic acid), HBPD (N,N'-Bis(2-hydroxybenzyl)propylene-1,3-diamine-N,N-diacetic acid), 2,3-dihydroxybebezoic acid, DTPA [diethyltriamine pentaacetic acid], and iron chelators produced by bacterial siderophores. The term "lipid carrier," as used herein, includes a carrier comprising lipid molecules, e.g., liposomes. Liposomes are spherical structures made of materials having a high lipid content, e.g., surfactants or phospholipids. The lipids of spherical vesicles are organized in the form of lipid bilayers. The lipid bilayers encapsulate an aqueous or oil-filled volume which is either interspersed between multiple onion-like shells of lipid bilayers (forming multilamellar lipid vesicles) or the aqueous volume is contained within an amorphous central cavity. The most commonly

known lipid vesicles having an amorphous central cavity filled with aqueous medium are the unilamellar lipid vesicles.

The present invention provides delivery systems for iron chelators. A hydrophilic drug, for example, Desferrioxamine, can be combined with a lipid carrier.

5 The iron chelator can be coated with the lipids of the lipid carrier, e.g., iron chelators associated, intercalated, or attached to the surface of lipid molecules or to the lamellae of a liposome. Alternatively, the iron chelator can be encapsulated within the lipid carrier, e.g., iron chelators encapsulated within the central cavity of a liposome, intercalated within the lamellae of a liposome, or located between lamellae of the

10 liposome. Other examples of lipid carrier formulations for the present iron chelator delivery system include, for example, the following: systems used with amphotericin B, which involve complexing the active ingredient with phospholipids as used in the formulation for ABELCET® (The Liposome Company, Inc., Princeton New Jersey); cholesteryl sulfate complexes for injection similar to the formulation used for

15 AMPHOTEC® (Sequus Pharmaceuticals, Menlo Park, California) which comprises a sterile, pyrogen-free, lyophilized powder for reconstitution and intravenous administration, e.g., a formulation comprising a complex of Desferrioxamine and cholesteryl sulfate (upon reconstitution a colloidal dispersion of microscopic disc-shaped particles result); and a single bilayer liposomal drug delivery system such as

20 used with AmBisome® (Nexstar Pharmaceuticals, Boulder Colorado (taken over by Gilead Sciences Foster City, California)), wherein single bilayer liposomes are used. The drug becomes active when the lipid carrier, e.g., the liposome, fuses with cells to release its content.

There are many advantages for using the iron chelator delivery system of

25 the present invention. The kidneys rapidly excrete small, hydrophilic iron chelator molecules, for example, Desferrioxamine. The iron chelator delivery system of the present invention, e.g., iron chelators associated with lipid carriers, e.g., liposomes, would not be filtered by the renal glomerulus. Even small unilamellar liposomes are about 10nm in diameter, well above the filtration limit of the glomerulus. Therefore, the

30 biological half-life of the iron chelator is extended when encapsulated within a liposome or associated with a lipid carrier.

Lipid carriers such as liposomes have been used for drug delivery for years with only limited success. A major problem has been targeting the liposomes to the proper organ or tissue. Attempts have been made to target chemotherapeutic agents

35 which are encapsulated in liposomes to tumor cells, using a guidance system such as a tumor-specific antibody fragment embedded in the lipid bilayer. A problem associated with this technique is that many, if not most of the liposomes, are trapped by the liver and engulfed by the Kupfer cells in this organ. However, the liver is the major target

organ in iron-overload, being the body's primary iron storage site. Therefore, reticuloendothelial (RE) cells engulf the iron chelator delivery system of the present invention, placing them in a position to intercept iron as it is released from degraded erythrocytes. Further, cardiac failure as a result of iron-overload is of great clinical importance. The heart can be targeted for delivery of Applicant's iron chelator delivery system, for example, Desferrioxamine combined with a lipid carrier by placing a cationic or anionic charge on the lipid carrier or by attaching antibodies specific to cardiac, vascular, endothelial, and matrix proteins to the lipid carrier. Also, the lipid carrier can be tagged with cardiac, liver, or tissue imaging labels. Cardiac proteins useful for targeting the heart include, for example, cardiac specific proteins such as myosin, troponin, and light chain myosin. Antibodies specific to endothelial, vascular smooth muscle cells, and matrix proteins can also be attached to the lipid carrier of the present iron chelator delivery system. In addition, selected delivery routes and an increase in the half life of the liposomes can enhance delivery to the heart.

Mammals, e.g., humans or animals, with transfusional iron-overload receive blood on a three to six week schedule, depending on the severity of their anemia. With the iron chelator delivery system of the present invention, the iron chelator delivery system, for example, Desferrioxamine combined with a lipid carrier, e.g., a liposome, would be infused at the end of the transfusion session and infused periodically. This would allow significant improvement over the current daily use of an infusion pump for 12-16 hours intervals. For example, thirty grams of Desferrioxamine encapsulated by liposomes would supply enough chelator for a month. This approach would be simpler than any available to date, including oral deferipone. Prior to administrations, the iron chelator delivery system can be dissolved in a pharmaceutically acceptable carrier. As used herein "pharmaceutically acceptable carrier" includes any and all excipients, solvents, dispersion media, coatings, antibacterial and antifungal agents, toxicity agents, buffering agents, absorption delaying or enhancing agents, surfactants, and micelle forming agents, lipids, liposomes, and liquid complex forming agents, stabilizing agents, and the like. The use of such media and agents for pharmaceutically active substance as known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

Conventional Desferrioxamine therapy costs currently between about \$12,000 and \$15,000 per year. Much of the expense involves the portable home infusion pumps, associated equipment (e.g., sterile infusion tubing and needles), and home nursing visits. These costs are eliminated by the current iron chelator delivery system. The iron chelator delivery system of the present invention can make iron

chelation therapy available to underdeveloped countries that lack the financial resources needed for conventional Desferrioxamine chelation programs.

The iron chelator delivery system of the present invention eliminates the problems of local reactions associated with the subcutaneous administration of some iron chelators. For example, the chelator diethyltriamine pentaacetic acid (DTPA) was used to treat iron-overload early on, but caused severe local reactions that led to discontinuation of use. Intravenous administration, although more efficacious, is associated with the risk of infection, atrial thrombosis, and subclavical thrombosis. The indwelling catheter often curls into the atrium causing cardiac irritation and thrombosis. Right atrial thrombosis can lead to life threatening pulmonary embolism. The iron chelator delivery system as described herein would negate these problems.

The iron chelator delivery system of the present invention uses the chelator more efficiently. Compared to conventional therapy, lower amounts of chelator are required with Applicant's iron chelator delivery system which combines iron chelators with lipid carriers, e.g., liposomes. For example, by concentrating the iron chelator delivery system in the reticuloendothelial (RE) cells initially, the chelator has the greatest possible chance of intercepting iron. Therefore, lower dosages (and cost) of the drug are attained.

The lipid carriers of the present invention, e.g., liposomes, have a number of advantages over currently used options such as red cell ghosts as encapsulation agents. Liposomes are manufactured, while red cells ghosts are harvested. Therefore, supply is not an issue. Use of red cell ghosts entails some exposure, even slight, to biological pathogens. Finally, lipid carriers, e.g., liposomes, avoid alloimmunization issues because they lack red cell antigens (Rose, W. et al. (1990) Blood 76:1431-7).

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

Example 1: Analysis of Iron-Overloaded Animals Used in the Following Experiments

Eighty Hartley Guinea Pigs weighing 350-400 grams of both sexes were purchased. Iron dextran was administered at 1.5 g Fe/kg body weight intraperitoneal three times a week for four weeks. Animals were divided into groups to receive iron chelator without lipid carrier and chelator with lipid carrier as described herein. Desferrioxamine was given intramuscularly twice a day daily for a total concentration of

20 mg/kg daily for five days a week for four weeks. Intravenous lipid associated chelator was given intravenously three times a week at 0.6 mg.

Iron-overloaded animals were sacrificed and various organs were analyzed. The pattern of changes in the organs of all the iron-overloaded animals studied were similar with slight differences in severity. In the livers, there was a mild to moderate Kupffer cell hyperplasia in hepatic sinusoids. The Kupffer cells were distended with black globular material (Hematoxylin and Eosin) that was iron positive (Perl's iron stain). The Kupffer cells were clumped (granulomas) in those animals having moderate hyperplasia. The clumping was random and often associated with degeneration, necrosis, and replacement of hepatocytes. Rarely, lymphocytes were admixed. Hepatocytes also contained small amounts of iron positive material detected with special stains.

The hearts were characterized by cellular infiltration of macrophages containing iron. These cells were present in the epicardium and myocardium (perivascularly and in the interstitium) and in the adipose tissue at the base of the heart and perivascularly around the pulmonary vessels.

This model of iron overload therefore reflects the human condition and is suitable for testing the iron chelator delivery system described herein. Guinea pigs have been treated with the non-lipid associated chelator (8 mg/day 5 days per week) and the lipid-associated chelator (8 mg/day three days per week). In two animals, after only three injections of the lipid associated chelator, the color of the eyes and ears were restored to that seen in non-overloaded Guinea Pigs.

Example 2: Preparation of an Iron Chelator Delivery System

Iron chelator delivery systems comprising an iron chelator and a lipid carrier, e.g., a liposome, can be prepared using the following standard procedures. General methods for preparing liposomes for use in the present iron chelator delivery systems are provided in Cortesi R., Esposito E., Gambarin S., Telloli P., Menegatti E., Nastruzzi C. (1999) "Preparation of liposomes by reverse-phase evaporation using alternative solvents," *Journal of Microencapsulation* 16(2):251-256; Buboltz J.T., Feigenson G.W., (1999) "A novel strategy for the preparation of liposomes: rapid solvent exchange," *Biochimica et Biophysica Acta* 1417(2):232-245; Bandyopadhyay P., Kren B.T., Ma X., Steer C.J., (1998) "Enhanced gene transfer into HuH-7 cells and primary rat hepatocytes using targeted liposomes and polyethylenimine," *Biotechniques* 25(2):282-284, 286-292; Puu G., Gustafson I., (1997) "Planar lipid bilayers on solid supports from liposomes-factors of importance for kinetics and stability," *Biochimica et Biophysica Acta* 1327 (2):149-161; Weiner A., (1994) "Liposomes for protein delivery:

selecting manufacture and development processes,” Immunomethods 4(3):201-209), the contents of which are incorporated herein by reference.

To determine the λ_{\max} value for Desferrioxamine, a solution of Desferrioxamine Mesylate in methanol is prepared and absorbance spectra determined using a Perkin Elmer Lambda 3B UV/VIS Spectrophotometer (Figures 1 and 2).

To obtain a calibration curve (Figures 3 and 4), 10mg Desferrioxamine Mesylate is dissolved in 1200 ml of methanol and a stock solution generated. 0.1 ml, 0.25 ml, 0.5 ml, 0.75 ml, 1.0 ml, 1.5 ml, and 2.0 ml of this stock solution is transferred to separate volumetric flasks to form a final volume of 10 ml with the addition of methanol. Spectrophotometric absorbance for each flask is read at 204nm.

Method 1: Liposomes are prepared by the reverse phase evaporation technique (Tabak, A. et al. (1994) J. Pharm. Pharmacol. 46:789-796). Phosphatidyl choline (PC) 0.5 mg and 4.5 mg cholesterol (Ch) are dissolved in 4 ml of chloroform (CHCl_3). The aqueous phase containing 12 mg Desferrioxamine Mesylate in 2 ml of saline is added to the above organic phase. The two phases are vortexed for 4 minutes for formation without emulsion. Liposomes are obtained by evaporating the organic phase under a partial vacuum produced by a water aspirator on a rotary evaporator at 20° C-30°C. The resulting liposomes are extruded through 1.0, 0.6, 0.4, and 0.2 μm Nucleopore polycarbonate membrane filters sequentially (membrane filter pore sizes may be varied to obtain liposomes of larger sizes). The nonliposomal Desferrioxamine is removed by using the Sephadex G-50 spin-column centrifugation method. Briefly, the bottom of a 3 ml disposable syringe (0.5 cm internal diameter) is plugged with clean, silanized glass wool. A Sephadex G-50 slurry is loaded into the syringe – column. The syringe is placed into a 15 ml polypropylene tube and spun for 3 minutes at 12500 rpm in order to pack the column. A 0.25 ml sample is then loaded onto the column and centrifuged for 3 minutes at a speed of 1500 rpm. Twenty-five μl of eluant is transferred to a volumetric flask and made up to a final volume of 10 ml with methanol. The flask is then shaken to extract intraliposomal Desferrioxamine. Spectrophotometric absorbance of the solution is determined at 204 nm.

Method 2: A mixture of phosphatidyl choline (PC) and cholesterol (16 mg PC: 8 mg Ch) in CHCl_3 is dried under a stream of nitrogen and vacuum desiccated for 2 hours to remove any remaining traces of the solvent. The lipid is then hydrated by addition of 14 mg of Desferrioxamine in 4 ml of saline. Saline is passed through a Chelex column. The mixture is vortexed for 5 minutes. The resulting liposomes are extruded through 1.0, 0.6, 0.4, and 0.2 μm Nucleopore polycarbonate membrane filters, sequentially (membrane pore sizes may be varied to obtain liposomes

of larger sizes). Liposomes are dialyzed overnight against saline at 4° C to remove non-encapsulated Desferrioxamine. Intraliposomal Desferrioxamine is assayed by dissolving an aliquot of the suspension in an equal volume of Triton X 100, adding FeCl₃ in 0.1 M HCl and measuring the absorbence of the iron-Desferrioxamine complex at 428 nm

5 (Young, S.P., et al. (1979) Brit. J. Haem. 41:357-363).

Example 3: Preparation of Radiolabeled Liposomes with ¹¹¹In

Desferrioxamine is prepared by activation of the carboxylic groups of Desferrioxamine with carbodiimide at pH 4.0 using a method similar to that described

10 by Khaw et al. (Khaw, B.A. et al. (1991) J. Nucl. Med. 32:1742-1751). 8.33 X 10⁻⁵ mM DTPLA in 250 µl H₂O (pH adjusted to 4.0) is mixed with 14-fold molar excess of N-hydroxy sulfosuccinimide (NHSS) and 400 µg/40 µl 1-ethyl-3-(3-dimethylamino-propyl) carbodiimide (EDC) is added. Another aliquot of NHSS is added. The reaction is allowed to proceed for 5 minutes at room temperature. The activated

15 Desferrioxamine is added then directly into 1 mg of phosphatidylethanolamine in 0.1 M borate buffer at pH 8.3. At this pH, there is unidirectional coupling of the carboxyl group of the activated Desferrioxamine to the amino group of the phosphatidyl ethanolamine. Free Desferrioxamine is separated from Desferrioxamine-PE by lypophalization of the mixture and separation of the Desferrioxamine-PE by extraction

20 with chloroform. Desferrioxamine will not be soluble, but Desferrioxamine-PE will be soluble. The chloroform is evaporated to obtain Desferrioxamine-PE. Aliquots of Desferrioxamine-PE is solubilized in 0.5 M citrate pH 5.0 to which 1-2 mCi of ¹¹¹In is added. The ¹¹¹In-Desferrioxamine-PE is added to cholesterol and lecithin to prepare liposomes as described below.

25

Example 4: Glycosylated and Mannosylated Lipid Carriers, e.g., Liposomes

Glycosylated liposomes are prepared by linking liposomes to a hydrophobic anchor N-glutaryl phosphatidyl ethanolamine (NGPE), which have been previously derivatized with the sugar residues. The carboxylic groups of NGPE are first

30 activated by water soluble 1-ethyl-3 (3-dimethylaminopropyl) carbodiimide (EDC) to which ethylenediamine is then added to provide free amino groups. Thiolated galactose and mannose residues are then incorporated onto the amino groups of the ethylenediamine by the method of Lee et al. (Lee, Y.C. et al. (1976) Biochem. 15:3956-3963).

35

Example 5: Iron Chelator and Galactosylated and Mannosylated Lipid Carriers, e.g., Liposomes

N-glutaryl phosphatidyl ethanolamine (NGPE) (0.075 mg) is dissolved in 2-(N-morpholino) ethanesulfonic acid hemisodium salt (MES) buffer (0.016M octylglucoside in 50 mM MES). After addition of 0.6 mg 1-ethyl-3-(3-dimethyl amino propyl) carbodiimide (EDC) the resulting mixture is incubated at room temperature for 5 minutes. Ethylenediamine (100 μ M) is then added to the activated NGPE and the solution kept at pH 5.0 by the addition of 1 N HCl solution. The reaction is allowed to proceed overnight with stirring and the solution is dialyzed against distilled water. Sugar residues are incorporated onto the amino groups of ethylenediamine-coupled NGPE by using 2-imino-2-methoxyethyl 1-thiogalactoside or 1-thio-mannoside to obtain galactosylated and mannosylated NGPE.

Liposomes are prepared by reverse phase evaporation technique (Ashwell, G. (1974) Adv. Enzymol. 41:99-128) as previously described and outlined above. The aqueous phase containing Desferrioxamine Mesylate and either galactosylated or mannosylated NGPE are added to the solubilized lipids. The two phases are vortexed for formation without emulsion. Liposomes are obtained by evaporating the organic phase under a partial vacuum produced by a water aspirator on a rotary evaporator. The resulting liposomes are extruded through Nucleopore polycarbonate membrane filters. The nonliposomal Desferrioxamine will be removed by Sephadex G-50 spin-column centrifugation method as described above.

Example 6: Iron Chelator and Cationic Tagged Lipid Carriers, e.g., Liposomes

Positive (cationic) liposomes are prepared according to Caride and Zarel (Caride, W. et al. (1977) Science 1989:735-738). Cationic liposomes are prepared using lecithin, cholesterol and stearylamine (80:10:10 molar ratios). Incorporation of Desferrioxamine into the cationic liposomes is as previously described. Trace amounts of In-111 DTPA-PA are added to radiolabel liposomes.

Example 7: Preparation of Fluorescein Desferrioxamine

An aliquot of 120 mg of Desferrioxamine and 233.64 mg of fluorescein isothiocyanate (FITC) are mixed in 50ml of 0.1 M Na carbonate pH 9.5. A molar ratio of 3:1 of FITC: Desferrioxamine is used as described in Hentz et al. (1997) Anal. Chem. 69:4994-5000. The reaction mixture is wrapped in aluminum foil to avoid light exposure and allowed to react for at least 24 hours at room temperature. To separate fluorescent- Desferrioxamine from free fluorescein and free Desferrioxamine, the mixture is electrophoresed over a P-2 column (Bio-Rad) equilibrated in saline and samples are eluted in 2 ml fractions. The fluorescein- Desferrioxamine is eluted in the void volume of the column. The first peak is collected and used for preparing fluorescein-Desferrioxamine, Desferrioxamine, and rhodamine-liposomes.

Example 8: Preparation of Rhodamine Labeled Liposomes Containing Unlabeled Desferrioxamine and Rhodamine Labeled Liposomes Containing Fluorescein Labeled Desferrioxamine

5 An aliquot of 75 mg 1- α -phosphatidyl choline (20 mg/ml chloroform, Avanti Lipids) is added to a 50 ml round bottom flask. Forty-five mg of cholesterol (cholestin-3 β -ol) is added to the solution of phosphatidyl choline. To this is added 373 μ l of 2 mg/ml 1,2 dipalmitoyl-sn-glycero-3-phosphaethanolamine-N-(lissamine rhodamine β sulfonyl) (Avanti Lipids). The mixture is stirred by vortexing until

10 thoroughly mixed. The mixture is then evaporated with a rotary evaporator set at 42°C for 1 hour at 160 rpm. One hundred and twenty mg of Desferrioxamine is dissolved in 20 ml of physiological saline and then added to the evaporated lipids in the rotary evaporation flask. To this mixture is added 10% (vol/vol) of fluorescein-Desferrioxamine (approximately 2 ml). The mixture is stirred vigorously with a

15 vortexed mixer. The unilamella liposomes are chromatographed using multiple columns of PD-10 (Pharmacia, Sephadex-G25) for column centrifugation as described. The liposomes that are eluted in the eluate are pooled and extruded at least 5 times each serially through 0.8, 0.45, and 0.22 micron diameter nucleopore filters. A total of 120 mg of lipids is collected in a total of 20 ml of physiological saline. Assuming a loss of

20 20% in the column centrifugation, the final lipid concentration in the 20 ml sample is approximately 96 mg/ml of Desferrioxamine and 9.6 mg of fluorescein-Desferrioxamine.

To prepare rhodamine liposomes with non-fluorescent Desferrioxamine, the fluorescein-Desferrioxamine is not included in the above formulation and the

25 remainder of the protocol remains the same.

Example 9: Targeted Delivery of Iron Chelator System

Targeted delivery of iron chelator delivery systems comprising, for example, Desferrioxamine and liposomes, can be shown by comparing the delivery of

30 the liposomes in normal animals with the delivery of the liposomes in animals with iron-overload. Radiolabeled liposomes can be used to quantify the localization into targeted tissues and, for example, the effect on heart function in particular arrhythmogenesis and contractile dysfunction can be evaluated.

The Heart: Iron chelator delivery systems comprising, for example,

35 Desferrioxamine and liposomes, for targeting the heart include placing a cationic or anionic charge groups on the liposomes which might adversely affect the conduction system of the heart. Injection of the Desferrioxamine and liposome system can be done intra-venously. Caride and Zaret in 1977 (Caride, W.J. (1977) Science 198:735-738)

showed that positively charged liposomes can be targeted specifically for acute myocardial ischemia or infarction in an experimental model. When uptake was compared to regional myocardial blood flow, it was observed to have an inverse correlation to the liposome distribution. Negatively charged (anionic) liposomes, however, distributed relative to blood flow. Since there is a myocardial blood flow, it was observed to have an inverse correlation to the liposome distribution. Since there is myocardial involvement in iron toxicity, there is possibility of damage to the myocardium. Therefore, cationic or anionic liposomes will also localize preferentially in iron-overloaded myocardium. Another approach would be to tag the iron chelator delivery system with antibodies targeted to cardiac specific endothelial cells, vascular smooth muscle cells, matrix proteins, receptors, and/or myocardial cells.

The Liver: Carbohydrate receptors in the liver, such as asialoglycoprotein receptors on the hepatocytes and mannose receptors on Kupfer cells and endothelial cells, allow selective targeting of this organ (Ashwell, G. (1974) Adv. Enzymol. 41:99-128), (Ashwell, G. (1982) Ann. Rev. Biochem. 51:531-554), (Fallon, R.J. (1989) Adv. Drug Del. Rev. 4:49-63), (Gordon, S. (1989) Adv. Drug Del. Rev. 4:27-47), (Meijer, D.K.F. (1989) Phar. Res. 6:105-118) by the Desferrioxamine-encapsulated liposomes. Liposomes can be utilized as carrier molecules because of their low immunogenicity, their relative biocompatibility, and their drug encapsulation efficiency. With glycosylated liposomes the targeting efficiency to the liver is increased.

Example 10: Determination of the Toxicity of The Iron Chelator Delivery System

Toxicity in the administration of the iron chelator delivery system, e.g., Desferrioxamine combined with a lipid carrier, e.g., a liposome, can be tested in animal models. For example, two species of normal animals and iron-overloaded animals (e.g., the guinea pig) can be compared. One group of animals receives the delivery system comprising Desferrioxamine and a lipid carrier, e.g., Desferrioxamine-encapsulated within a liposome, and another group receives similar liposomes without the Desferrioxamine. Animals are monitored for cardiac arrhythmias via electrocardiogram once a week. Heart function is determined with echocardiography at the same time that electrocardiograph recordings are made at one and at 24 hour intervals post-injection followed by weekly monitoring for up to 6 weeks. Full necropsy is performed with special attention to harvesting the liver and heart. In humans, the infusion of the iron chelator delivery system, e.g., Desferrioxamine combined with a lipid carrier, e.g., Desferrioxamine encapsulated within a liposome, would take between 20 minutes and one hour. Repeat treatments typically would occur over a 3-6 week interval. Therefore cardiac and liver functions are monitored in the animals after one hour, 24 hours, and at weekly intervals up to ten weeks.

Local skin irritation at the infusion site is a complaint of the current formulation of Desferrioxamine. Irritation should not occur with the use of the present iron chelator delivery system, e.g., Desferrioxamine encapsulated within a liposome, because liposomes are given intravenously. Animal body weight, serial hematocrits, blood iron levels, and ferritin plasma levels are determined on individual animals. Blood samples are analyzed with full differential including creatinine and complete CBC to check for agranulocytosis as seen with Deferipone. Analysis of liver enzymes including SGOT, SGPT, alkaline phosphatase, BUN, is performed once a week. Urine excretion or iron is measured in iron-overload animals by atomic absorption spectroscopy.

Dosing range, safety, and toxicity studies in normal animals, as well as, quantification of any toxicity seen in the liver, heart, gastrointestinal tract, kidney, and at the injection site is important. At the time of full body necropsy tissue weights and gross appearance is recorded. The heart and liver are stored in 10% phosphate buffered formalin. Other body organs are also stored in 10% buffered formalin for future light microscopy, or special staining. Prussian blue iron stain is applied to liver and heart samples. Quantitative analysis using a point counting technique and semiquantitative technique using a scale of ± 4 can be used. All measurements should be made by a single observer.

Flame photometry atomic absorption spectroscopy measurements are carried out on fresh or lyophilized liver and heart with either normal iron depot or iron-overload with or without the present iron chelator delivery system, e.g., Desferrioxamine encapsulated within a liposome, administered. Normal animals treated with Applicants' iron chelator delivery system, e.g., Desferrioxamine encapsulated within a liposome, and non-treated normal animals are compared for possible reduction in normal iron levels. Iron deficiency can in itself result in cardiomyopathy. Similar studies would be applied to Mongolian gerbils and rats with iron-overload and thalassemic rats.

Example 11: Determination and Comparison of the Biological Half-Life of Iron Chelator Delivery Systems

Iron-overloaded rats are given iron chelator delivery systems comprising, for example, Desferrioxamine and liposomes. Iron blood levels are measured before and after injection. Pharmacokinetic studies are performed in both iron-overloaded and normal animals with radiolabeled liposomes. One group of animals also receives the standard formulation of Desferrioxamine injected over a 24 hour period via a minipump similar to the standard procedure for humans receiving the compound. The localization of Desferrioxamine will be radiolabeled for imaging and quantification of organ localization.

Imaging: Male and female rats age 10-12 weeks are injected intravenously with 100 μCi ^{111}In -Desferrioxamine liposomes (4 $\mu\text{Ci}/\mu\text{mol}$ lipid, fluid liposome type) or 100 μCi free ^{111}In -Desferrioxamine. The amount of Desferrioxamine to be injected is determined in experiments without radiolabeling using fluorescent
 5 labels. Rats are anesthetized (halothane-nitrous oxide-oxygen) and placed prone on a single head camera equipped with a parallel hole, medium energy collimator to image the entire body with special focus on liver, kidney, brain, and heart. Symmetric 20 percent windows are focused for both 173 and 247 KeV energy peaks. Images (100,000 counts/image) are obtained and stored in a 256x256 matrix. The scintigraphic results
 10 are analyzed by drawing regions of interest. Images are obtained at 0.5, 1, 2, 4, 6, 24, 48, 72, 96, and 144 hours after injection of ^{111}In -Desferrioxamine-liposomes and free drug (five rats per time point).

The amount of liposome-entrapped Desferrioxamine was found to be highest in the liver and had a longer biological half-life than the iron chelator alone
 15 (Figures 6-18).

Pharmacokinetics/Biological Half-Life Determinations: Pharmacokinetic studies are performed in both iron overloaded and normal animals with radiolabeled liposomes. Pharmacokinetic parameters are calculated using the curve fitting program KINFIT. Values of ^{111}In -Desferrioxamine in blood versus time are fitted by a model for
 20 extravascular administration and two compartments without lag time. The half-life of blood levels of liposome-entrapped Desferrioxamine is determined.

Example 12: Toxicity Study in Normal Animals

Safety studies were performed in six 200 gram female rats. Three rats
 25 were injected via tail vein with 2 ml of iron chelator delivery systems comprising, for example, Desferrioxamine and liposomes, (10 mg). Three additional animals received the liposomes without Desferrioxamine. No toxicity or morbidity was noted up to ten days post-injection. There was no irritation noted at the injection site.

Blood Chemistry Values Before and After Liposome Infusion (with and
 30 without Desferrioxamine) were not significantly different (Figure 5).

Rats were anesthetized and allowed to breathe spontaneously during fluid filled catheter recordings of LV pressure. Closed chest LV systolic pressures were 180 \pm 4 mm Hg before intravenous injection of the Desferrioxamine-encapsulated liposomes versus 181 \pm 4 after injection. At one hour after injection the LV systolic
 35 pressure was 175 \pm 5 mm Hg ($p > 0.05$).

Simple semiquantitative analysis of tissues from animals receiving the liposomes without Desferrioxamine and liposomes with Desferrioxamine (i.e., liver and heart) stained for iron revealed no differences by light microscopic examination.

Example 13: Histological Findings of Rats Injected with Rhodamine Labeled Liposomes Containing Fluorescein Labeled Desferrioxamine

Forty-two rats were injected with 2 cc of rhodamine labeled liposomes containing 0.6 mg/ml of Desferrioxamine labeled with fluorescein. Heart, liver, brain, and both kidneys were harvested. Tissues were frozen for later fluorescent imaging or stored in buffered formalin, sectioned, and stained with hematoxylin and eosin and counter stained with Masson's trichrome. Histological reading by a board certified laboratory veterinary pathologist reported no remarkable change. No morbidity or toxicity was noted up to ten days post-injection. There was no irritation noted at the injection site. An additional group of animals were euthanized at 30 minutes, 1 hour, and 24 hours post-injection of Desferrioxamine labeled with fluorescein entrapped in liposomes labeled with rhodamine. Histological analysis showed fluorescent labeling in the liver of liposomes and of Desferrioxamine in the Kupffer cells at the 30 minute time point. Discrete localization was seen up to two hours post-injection. The 24 hour time point demonstrated homogenous fluorescence due to re-distribution of the lipids in the liposomes and Desferrioxamine. There was no detected organ damage. No signs of toxicity were noted in any of the experiments at the organ level or clinical presentation. The brain was examined for histological changes as lipids can easily cross the blood brain barrier. No indication was found of liposome distribution in the brain, heart, or kidney.

Example 14: Biodistribution of the Iron Chelator Delivery System

Rats are injected intravenously with 10 μCi ^{111}In -Desferrioxamine liposomes or 10 μCi free ^{111}In -Desferrioxamine at a total dose of 25 μmol Desferrioxamine targeted (injection volume 0.5 ml). At predefined time points after injection, rats receive an overdose of pentobarbitol and blood is obtained by cardiac puncture. The following tissues are dissected: injection site; muscle; lung; spleen; kidney; liver; heart; and brain. These tissues are also weighed and the activity measured in a shielded well-type gamma counter. To correct for physical decay and to calculate uptake in each organ as a fraction of the injected dose, aliquots of the injected dose are counted simultaneously.

With reference to Figures 6-18, following the injection of radiolabeled Desferrioxamine, radioactivity is observed in the kidney and the spleen. Lung, spleen, blood, skeletal muscle, brain, heart, and liver are also radioactive. No radioactivity is observed in the brain, indicating that the iron chelator delivery system does not cross the blood brain barrier. As the radioactivity in the blood drops over time, the liver

localization of the radioactive iron chelator delivery system in the liver increases. Subsequently, the chelated Desferrioxamine is removed from the liver.

Radiolabeled chelator without liposomes was quickly eliminated from the animals and did not accumulate in the tissues, especially the liver and tracked blood
5 distribution.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to specific embodiments of the invention
10 described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194